

ST. GABRIEL LABORATORIES

Fast Acting

BurnOut II

**NEW &
IMPROVED
FORMULA**

WEED & GRASS KILLER

CONCENTRATE

For Non-Selective Control of
Herbaceous Broadleaf and
Grass Weeds.

Active Ingredients:

Clove Oil 12.00%

Sodium Lauryl Sulfate ... 8.00%


Other Ingredients:

Vinegar, Lecithin, Water,

Citric Acid, Mineral Oil.

Total Other 80.00%

Total.....100%



KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel booklet for additional precautionary statements

Net Contents: 2.5 US Gallons

ST. GABRIEL LABORATORIES

Fast Acting **BurnOut™** NEW & IMPROVED FORMULA **WEED & GRASS KILLER CONCENTRATE**

St. Gabriel Laboratories

14044 Litchfield Drive
Orange, Virginia 22960
1-800-801-0061
www.milkyspore.com

To reorder call: 800-801-0061

KEEP OUT OF REACH OF CHILDREN **CAUTION** FIRST AID

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND ANIMALS

CAUTION: Corrosive, causes irreversible eye damage. Harmful if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield when handling. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTION EQUIPMENT (PPE): Applicators and other handlers must wear: Appropriate protective eyewear, such as goggles or face shields, long-sleeved shirt and long pants, waterproof gloves and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for PPE, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS: Avoid spraying directly into water.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep from freezing. Store only in original tightly sealed container and out of reach of children.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container (bottle). Rinse thoroughly before discarding in trash.

CONDITIONS OF SALE

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffective or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide-resistant weed preparations or the use or application of the product contrary to label instructions, all of which are beyond the control of St. Gabriel Laboratories. ALL SUCH RISKS WILL BE ASSUMED BY THE USER. St. Gabriel Laboratories shall not be responsible for losses or damages resulting from the use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label. St. Gabriel Laboratories warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. ST. GABRIEL LABORATORIES DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. St. Gabriel Laboratories EXCLUSIVE REMEDY AND St. Gabriel Laboratories EXCLUSIVE LIABILITY SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THIS PRODUCT. In no case shall St. Gabriel Laboratories or the seller be liable for consequential, special, or indirect damages resulting from the use of this product. The foregoing is a condition of sale by St. Gabriel Laboratories and is accepted as such by the buyer.

DIRECTIONS FOR USE **SHAKE WELL BEFORE USING**

BurnOut Weed & Grass Killer-Ready To Use is a foliar active weed control product which contains clove oil as the active ingredient and extracts from vinegar as one of the inert. It is non-selective to both broadleaf and grassy weeds and is recommended for areas such as borders, driveways, sidewalks, around mature tree bases, around buildings, in fence rows, in greenhouses, school grounds, and around flower beds.

BurnOut Weed & Grass Killer is fast acting. Foliar contact results in rapid control of annual weeds and grasses, and top growth reduction of herbaceous perennial weeds and grasses (See Tables below). Re-treatment is required for control of established perennial weeds.

SHAKE WELL BEFORE MIXING

Fill the spray tank 1/2 full with water and start agitation system if applicable. Add the desired amount of BurnOut concentrate at a rate of (3:1), 3 parts water to 1 part BurnOut concentrate, or (2:1), 2 parts water to 1 part BurnOut concentrate.

MIXING RATIO GUIDE:

For young tender weeds and grasses (3:1) mixing ratio can be sufficient to kill target plants. For older more established weeds and grasses a (2:1) mixing ratio may be required to kill target plants.

Once in suspension, occasional agitation is recommended for best results.

BROADCAST APPLICATIONS: Spray BurnOut to run-off using tank or backpack sprayers, typically 35-75 gallons of spray solution per acre.

BAND APPLICATION: Using trigger sprayers, cover desired weeds and grasses, as needed, typically 25-50 gallons of spray solution per acre.

Best results are achieved from spring/early summer applications to actively growing young weeds. Weeds that are mature, dormant, or hardened due to moisture stress are more tolerant of herbicide treatments. Only contacted vegetation will be affected. Rainfall within one hour of application.

Avoid application to reactive metals such as aluminum, tin, iron, or items such as fencing or lawn furniture in order to prevent staining, mottling, or otherwise interfering with finished metal surfaces. In case of contact, rinse the sprayed surfaces with water.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

NOTE: ALL CONTACTED VEGETATION WILL BE AFFECTED. AVOID CONTACT WITH DESIRABLE PLANTS. OVERSPRAY OR DRIFT WILL INJURE OR KILL CONTACTED VEGETATION.

TYPICAL WEEDS CONTROLLED

ANNUAL BROADLEAF WEEDS
Black medic
Chickweed
Cinquefoil (rough)
Cocklebur
Croutalaria
Curled dock
Fleabane spp.
Ivy Leaf Morning Glory
Lamb's-quarters
Mustard spp.
Oxalis spp.
Pigweed spp.
Ragweed spp.
Sicklepod
Vickweed
ANNUAL GRASSES
Crabgrass
Foxtail spp.

PERENNIAL BROADLEAF WEEDS
Cinquefoil (silvery)
Clover spp.
Dandelion
Plantain spp.
Toadflax
Turfed vetch
Wild carrot
PERENNIAL GRASSES
Bluegrass
Quack grass

OTHER
Algae
Liverwort
Moss

BARCODE