



# BugShooter

## GARDEN BUG SPRAY

### CONCENTRATE FORMULA

*For Crawling insects in and  
Around the Garden*

**DANGER**  
**KEEP OUT OF REACH**  
**OF CHILDREN**

See back panel for additional  
precautionary statements.

Active Ingredients:  
Clove Oil ..... 3%  
Peppermint Oil ..... 2.5%  
Sodium Lauryl Sulfate ..... 10%  
Inert Ingredients:  
(Mineral Oil (USP), Citric Acid,  
Lecithin, Water) ..... 84.5%  
Total ..... 100.00%



NET CONTENTS: 32 OZ. (946 ML)

# BugShooter

## GARDEN BUG SPRAY

### CONCENTRATE FORMULA

#### KEEP OUT OF REACH OF CHILDREN

#### FIRST AID

**IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation persists or occurs.

**IF SWALLOWED:** Call a doctor or get medical attention.

**IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation occurs or persists.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.


**STORAGE:** Keep from freezing. Store only in original tightly sealed container and out of reach of children.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse container (bottle). Rinse thoroughly before discarding in trash.

#### CONDITIONS OF SALE

Buyer assumes all responsibility for safety and use not in full accordance with directions for use.



#### St. Gabriel Organics

14044 Litchfield Dr., Orange, VA 22960

1-800-801-0061 • [www.StGabrielOrganics.com](http://www.StGabrielOrganics.com)

PEEL HERE  
TO OPEN LABEL

This panel blank / affixes to container

## DIRECTIONS FOR USE

**USE THIS PRODUCT IN A MANNER  
CONSISTENT WITH ITS LABELING.**

### SHAKE BEFORE USING

### MIXING INFORMATION:

#### IF TREATMENT AREA IS LESS THAN ONE ACRE:

Rate of application is 1 part of BugShooter Concentrate to 9 parts of water, or 14 ounces (27 tablespoons) of BugShooter Concentrate per gallon of water. Apply until foliage is wet or a maximum of two gallons per 1000 square feet.

Spray when insects first appear. Spray upper and lower leaf surfaces until run-off. Wet as many insects with spray as possible. Be sure to penetrate dense foliage. Treatment may be repeated weekly or as needed to control infestation. Avoid spraying plants in hot midday sun. When used outdoors, spray when air is still. Can be used up to day of harvest.

This product has been tested on many plants; however, it is wise to spray new plants individually and observe for 48 hours. Do not spray on open blooms and do not spray wilted plants.

Do not use in conjunction with other sprays such as spray-on liquid fertilizer and other pesticides. Do not spray on red mature fruit such as raspberries or strawberries. Do not spray at temperatures over 90°F. Early morning or evening spraying is best.

### USE ON VEGETABLES AND FRUITS INCLUDING:

Asparagus, beans, broccoli, brussel sprouts, cabbage, carrots, cauliflower, celery, collards, corn, cucumbers, eggplants, grapes, greens, kale, lettuce, peas, peppers, potatoes, radishes, spinach, squash, tomatoes, turnips, pumpkins, melons, zucchini, etc.

### USE ON HOUSEPLANTS, FLOWERS, TREES AND SHRUBS

including: (Test area on plant to check sensitivity) Asters, azaleas, begonias, boxwood, camellias, carnations, chrysanthemums, dahlias, dieffenbachia, dogwoods, ferns, fuchsias, geraniums, gladioli, hemlocks, hibiscus, holly, iris, ivy, jade plants, junipers, lilacs, maple, marigolds, oak, pepperomia, petunias, philodendron, rhododendrons, roses, rubber plants, salvia, snapdragons, spruces, tulips, wandering jews, etc., and **fruit trees**; such as: apples, bananas, cherries, figs, grapefruits, kiwi, lemons, limes, oranges, pears, plums, etc.

### USE TO CONTROL THE FOLLOWING INSECTS:

Ants (including fire ants), aphids, cabbage loopers, caterpillars, eastern and forest tent caterpillars, crossed-striped cabbageworms, cutworms, diamond-back moth larvae, earwigs, fireworms, flies, gnats, gypsy moth caterpillars, harlequin bugs, imported cabbageworms, lacebugs, leafhoppers, leaf rollers, loopers, mealy bugs, mites, moths, scales, snails/slugs, whiteflies, and mosquitoes, etc.

## PRECAUTIONARY STATEMENTS

### CAUTION:

May cause eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact, hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention if irritation occurs or persists. Wash thoroughly with soap and water after handling.