

Item#	Description	Case Pack	Size	Sh Wt.	UPC 037321	Case Cube
791	Dipel 150 Dust	12	1 lb.	14	007913	0.79
793	Dipel 150 Dust	12	4 lb. bags	50	007937	1.50

A biological insecticide that controls cabbage looper, imported cabbageworm, grape leaf folder and hornworm on crops listed on back panel.

Easy to use duster method.

DIPEL® 150 DUST is a biological insecticide intended for use against chewing larvae (caterpillars) of certain lepidopterous species. DIPEL® 150 DUST is recommended for use by ground or by air. Apply when pests are first noticed and repeat as necessary. Thorough plant coverage is necessary for control.

CROP	INSECT	APPLICATION RATE
Broccoli, Cabbage, Cauliflower, Collards, Celery, Melons, Potatoes, Spinach, and Cucumbers.	Diamondback Moth, Hornworms, Cabbage Looper, Melonworms, Webworm, Green Cloverworm	Apply 1/4 to 1/2 lb. to 435 sq. ft.
Broccoli, Cabbage, Cauliflower, Collards, and Turnip Greens	Loopers, Imported Cabbageworm, Hornworms	Apply 2 1/2 to 4 oz. to 435 sq. ft.
Grapes, Small Fruits, Cane and Bush Berries	Loopers, Omnivorous Leafrollers, Grape Leaf Folder, Grape Leaf Skeletonizer	Apply 1/4 to 1/2 lb. to 435 sq. ft.
Tobacco	Hornworm	Apply 2 to 4 oz. to 435 sq. ft.
Tomatoes, Peppers, Eggplants	Loopers, Hornworms, Tomato Fruitworm*, Variegated Cutworm, Salt Marsh Caterpillars, Armyworms**	Apply 2 to 4 oz. to 435 sq. ft.
*Apply weekly in a preventative program. **Dipel Dust may be used to suppress small worms (1st & 2nd instar) when populations are light and full ground coverage can be attained.		
Lawns	Sod Webworm	Apply 4 to 8 oz. per 100 sq. ft.
Flowers	Loopers, Tobacco Budworm, Omnivorous Looper, Omnivorous Leafroller, Diamondback Moth, Armyworms*, Ello Moth (Hornworm), Lo Moth, Oleander Moth	Apply 2 to 4 oz. to 435 sq. ft.

*Dipel Dust may be used to suppress small armyworms (1st & 2nd instar) when populations are light and full coverage ground sprays are applied.

PRECAUTIONARY STATEMENTS — CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

STATEMENT OF PRACTICAL TREATMENT: **If swallowed:** Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. **If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention. If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists. **If in eyes:** Flush eyes with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS: Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE — GENERAL CLASSIFICATION: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. **STORAGE:** Store in original container in a cool, dry place inaccessible to children and pets and away from heat and direct sunlight. Protect from freezing. Storage at temperatures above 90°F may impair effectiveness. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

EPA Est. No. 4-NY-1 (1), 769-GA-1 (2), 9198-OH-2 (3), 44616-MO-1 (4), 57062-OH-001 (5)

EPA Reg. No. 4-251

The last digit in the lot number indicates EPA establishment location.

ACTIVE INGREDIENT:

<i>Bacillus thuringiensis</i> subspecies <i>kurstaki</i> , Potency of 320 International Units per milligram (at least 0.5 billion viable spores per gram)*	0.064%
Lepidopteran Active Toxin	99.936%
INERT INGREDIENTS	100.000%
TOTAL	100.000%

*Equivalent to potency of 0.15 billion International Units per pound of this product.

Potency units should not be used to adjust use rates beyond those specified in the Directions for Use Section.

CAUTION KEEP OUT OF REACH OF CHILDREN

See back panel for additional precautionary statements.