

TRUSTED SINCE 1926

**BONIDE**

# Slug Magic™

Makes Slugs Disappear

## • Proven Snail and Slug killer

- For use around vegetables, fruit trees, citrus, berries, ornamentals, shrubs, flowers, trees, lawns, gardens and in greenhouses.
- Remains effective after rain or sprinkling



Can Be Used Around Pets and Wildlife

ACTIVE INGREDIENT:	
Iron phosphate.....	1.0%
INERT INGREDIENTS:.....	99.0%
TOTAL.....	100.0%

Keep out of reach of children

**CAUTION**

(See Back Panel for Additional Precautionary Statements)



## SLUG MAGIC™ Makes Slugs Disappear

- Can be used around pets and wildlife.
- May be used up to and including day of harvest.
- The bait is attractive to slugs and snails.
- Baits & Kills
- 1 lb. treats 1,000 sq.ft.

Made with Ferramol™ Slug and Snail Bait, a trademark of W. Neudorff GmbH KG

US Patent Number 5,437,870

EPA Est. No. 4-NY-1 EPA Reg. No. 67702-3-4

BONIDE PRODUCTS, INC.  
6301 Sutliff Road  
Oriskany, NY 13424

BN

Page 1

Directions for Use  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SLUG MAGIC™ Makes Slugs Disappear is a unique blend of an iron phosphate active ingredient, originating from soil, with slug and snail bait additives. It is used as an ingredient in fertilizers. The bait which is not ingested by snails and slugs will degrade and become a part of the soil.

The bait is attractive to slugs and snails and lures them from their hiding places and plants. Ingestion, even in small amounts, will cause them to cease feeding. This physiological effect to the bait gives immediate protection to the plants even though the slugs and snails may remain in the area. After eating the bait, the slugs and snails cease feeding, become less mobile and begin to die within three to six days. Dead slugs and snails may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the decrease of plant damage.

This product is effective against a wide variety of slugs and snails and will give protection to home lawns, gardens, greenhouses, outdoor ornamentals, vegetable gardens, fruit, berries, citrus and crop plants. The bait can be scattered on the lawn or on the soil around any vegetable crops, flowers, or fruit trees or bushes to be protected.

Page 2

**FIRST AID**  
If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**PRECAUTIONARY STATEMENTS**  
Hazards to Humans and Domestic Animals:  
Caution: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Environmental Hazards: For terrestrial uses. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

**NON-AGRICULTURAL USE REQUIREMENTS**  
The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.  
There is a restricted-entry interval of zero hours for this product.

Page 3

**STORAGE AND DISPOSAL**  
Do not contaminate water, food or feed by storage or disposal.  
Pesticide Storage: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.  
Container Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused products down any indoor or outdoor drain.

**HOW TO APPLY**  
The slug and snail bait granules should be scattered on the soil around or near the plants to be protected. For broadcast application standard broadcast spreaders may be used. For row application standard granular spreaders may be used. Apply bait evenly at approximately 1 lb. per 1,000 square feet (0.15 oz., or about 1 level teaspoon per square yard) and reapply as the bait is consumed or at least every two weeks. Do not place in piles. If the ground is dry, wet it before applying bait. The soil should be moist but with little or no standing water.  
Reapply as the bait is consumed or at least every two weeks. Apply more heavily if the infestation is severe, if the area is heavily watered or after long periods of heavy rain. See specific directions for different plant types and for inside greenhouses.

Page 4

### WHEN TO APPLY

Evening is the best time to apply the bait, as slugs and snails travel and feed mostly by night or early morning.

### WHERE TO APPLY

All likely areas of infestation should be treated, especially around the perimeter of garden plots because these pests travel into plant areas from daytime refuges. They favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery, and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra lawns, citrus groves, ivy patches, and other ground cover where they obtain shelter by day.

### OUTDOOR ORNAMENTALS

Scatter bait in a 6 inch circular band around the base of the plants to be protected at 0.15 oz., or 1 level teaspoon, per square yard. If plants are next to a grassy area, spread the bait between the ornamentals and the grass. Slugs and snails traveling to the plants will encounter the bait before reaching the plant. In these situations, scatter the bait around the perimeter of the plot at approximately 1 lb. per 1,000 square feet to intercept snails and slugs traveling to the plot.



### VEGETABLES

The bait can be used to protect any vegetables from slug and snail damage, including (but not limited to): artichokes, asparagus, beans, beets, blackeye peas, broccoli, Brussels sprouts, cabbage, cantaloupe, carrots, cauliflower, corn, cucumbers, eggplants, garlic, lettuce, onions, peas, peppers, potatoes, radishes, rutabagas, spinach, squash, Swiss chard, tomatoes, and turnips. Scatter the bait around the perimeter of the vegetable plot at approximately 1 lb. per 1,000 square feet to provide a protective "barrier" for slugs and snails entering the garden plot. If slugs or snails are inside the rows, then scatter the bait on the soil around the base of the plants and between the rows.

### FRUITS INCLUDING CITRUS

The bait can be used to protect fruits from slugs and snails, including (but not limited to): apples, avocados, apricots, cherries, grapes, melons, peaches, plums, citrus, pears. For seedlings spread the bait around the base of the stem. Apply at 0.15 oz., or 1 level teaspoon, per square yard, in a 6 inch circular band around the base of the plants to be protected. For older trees, spread the bait around the base of the tree to intercept slugs and snails traveling to the trunk. Apply the bait at approximately 1 lb. per 1,000 square feet for orchards using standard fertilizer granular spreaders.



### BERRIES

The bait can be used to protect berries from slugs and snails, including (but not limited to): strawberries, blackberries, blueberries, boysenberries, loganberries, raspberries. Spread the bait around the perimeter of the plot to intercept slugs and snails migrating toward the berries. Use a rate of approximately 1 lb. per 1,000 square feet and scatter by hand or with granular spreaders. If slugs and snails are already in the plots, then carefully spread bait between the furrows near the base of the plants. For small plots, treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively.

### GREENHOUSES

Where slugs or snails are a problem in greenhouses, scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply about 1/2 teaspoon per 9 inch pot.



### LAWNS

The bait can be used to protect lawns. When slugs or snails are detected, scatter the bait at a rate of approximately 1 lb. per 1,000 square feet. Scatter by hand or with granular spreader where the slugs or snails are observed.



\*The slugs and snails controlled by this product include (but are not limited to): Deroceras reticulatum (Field slug), Deroceras laeve (Smooth slug), Arion subfuscus (Dusky slug), Arion circumscriptus (Gray garden slug), Arion hortensis (Black field slug), Arion rufus (Large red slug), Arion ater (Large black slug), Limax flavus (Spotted garden slug), Helix spp., Helicella spp., and Cepaea spp.

\*The highly compressed pellets are easy to use, clean to handle and economical.

\*Can be used around domestic animals (pets) and wildlife.

WARRANTY - Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. Buyer assumes all risk of any such use. Seller makes no other warranties, either expressed or implied.

Page 5

Page 6

Page 7

Page 8